

# The Schizophrenia Proneness Instrument, Adult version (SPI-A)

Schultze-Lutter F, Addington J, Ruhrmann S, Klosterkötter J; Giovanni Fioriti Editore, Rome

## Summary of basic symptom criteria

|  | Basic Symptom Criteria |   |   |   |   |   | 7 | 8 | 9 |
|--|------------------------|---|---|---|---|---|---|---|---|
|  | 0                      | 1 | 2 | 3 | 4 | 5 |   |   |   |
| 1. Inability to divide attention (B1)                            |                        |   |   |   |   |   |   |   |   |
| 2. Disturbance of expressive speech (C5)                         |                        |   |   |   |   |   |   |   |   |
| 3. Disturbances of abstract thinking (O3)                        |                        |   |   |   |   |   |   |   |   |
| 4. Captivation of attention by details of the visual field (O7)  |                        |   |   |   |   |   |   |   |   |
| 5. Thought interference (C2)                                     |                        |   |   |   |   |   |   |   |   |
| 6. Thought blockages (C3)  |                        |   |   |   |   |   |   |   |   |
| 7. Disturbance of receptive speech (C4)                          |                        |   |   |   |   |   |   |   |   |
| 8. Thought pressure (D3)   |                        |   |   |   |   |   |   |   |   |
| 9. Unstable ideas of reference (D4)                              |                        |   |   |   |   |   |   |   |   |
| 10. Thought perseveration (O1)                                   |                        |   |   |   |   |   |   |   |   |
| 11. Decreased ability to discriminate between ideas and ... (O2) |                        |   |   |   |   |   |   |   |   |
| 12. Derealization (O8)   |                        |   |   |   |   |   |   |   |   |
| 13. Visual perception disturbances                               |                        |   |   |   |   |   |   |   |   |
| Changed perception of the face or body of others (D5)            |                        |   |   |   |   |   |   |   |   |
| Photopsia (F2)   |                        |   |   |   |   |   |   |   |   |
| Micropsia, macropsia (F3)  |                        |   |   |   |   |   |   |   |   |
| Near and tele-vision (O4.1)                                      |                        |   |   |   |   |   |   |   |   |
| Metamorphopsia (O4.2)  |                        |   |   |   |   |   |   |   |   |
| Changes in colour vision (O4.3)                                  |                        |   |   |   |   |   |   |   |   |
| Changed perception of patient's own face (O4.4)                  |                        |   |   |   |   |   |   |   |   |
| Pseudomovements of optic stimuli (O4.5)                          |                        |   |   |   |   |   |   |   |   |
| Diplopia, oblique vision (O4.6)                                  |                        |   |   |   |   |   |   |   |   |
| Disturbances of the estimation of distances or sizes (O4.7)      |                        |   |   |   |   |   |   |   |   |
| Disturb. of the perception of straight lines/contours (O4.8)     |                        |   |   |   |   |   |   |   |   |
| Maintenance of optic stimuli, 'visual echoes' (O4.9)             |                        |   |   |   |   |   |   |   |   |
| Partial seeing including tubular vision (O4.10)                  |                        |   |   |   |   |   |   |   |   |
| 14. Acoustic perception disturbances                             |                        |   |   |   |   |   |   |   |   |
| Changed intensity/quality of acoustic stimuli (F5)               |                        |   |   |   |   |   |   |   |   |
| Acoasms (O5.1)   |                        |   |   |   |   |   |   |   |   |
| Maintenance of acoustic stimuli, 'acoustic echoes' (O5.2)        |                        |   |   |   |   |   |   |   |   |

Rate maximum of frequency within the past 3 months:

- 0 = absent; (II) never present  
 1 = rare; (II) less than once in a month  
 2 = mild; (II) short periods about once in a month  
 3 = moderate; (II) several times in a month or weekly  
 4 = moderately severe; (II) several times in a week  
 5 = severe; (II) daily, periods of improvement possible  
 6 = extreme; (II) daily, but not necessarily continuously

7 = has always been present in same severity ('trait')

8 = definitely present, but severity unknown

9 = symptom definition questionably met

### **CRITERIA FOR COPER**

| A. Cognitive-perceptive basic symptoms  | Yes | No |
|---|-----|----|
| 1. Have at least 1 of the symptoms 5-14 been reported?  |     |    |
| 2. If 'Yes' to 1, has any of these symptoms first occurred more than 12 months ago?                         |     |    |
| 3. If 'Yes' to 1 and 2, are any of the symptoms with first occurrence more than 12 months ago scored '3-6'? |     |    |
| 4. 'Yes' to 1, 2 and 3? COPER criteria met.   |     |    |

### **CRITERIA FOR COGDIS**

| B. Cognitive disturbances  | Yes | No |
|--|-----|----|
| 1. Have at least 2 of the symptoms 1-9 been reported?            |     |    |
| 2. If 'Yes' to 1, are at least 2 of these symptoms scored '3-6'? |     |    |
| 3. 'Yes' to 1 and 2? COGDIS criteria met.                        |     |    |

Please check:

No     Yes

No     Yes

COPER ('Yes' to A4.)

COGDIS ('Yes' to B3.)